

approach 0. To do this we use our knowledge of the sine function. Because the sine of any number lies between  $-1$  and  $1$ , we can write

$$\boxed{4} \quad -1 \leq \sin \frac{1}{x} \leq 1$$

Any inequality remains true when multiplied by a positive number. We know that  $x^2 \geq 0$  for all  $x$  and so, multiplying each side of the inequalities in  $\boxed{4}$  by  $x^2$ , we get

$$-x^2 \leq x^2 \sin \frac{1}{x} \leq x^2$$

as illustrated by Figure 8. We know that

$$\lim_{x \rightarrow 0} x^2 = 0 \quad \text{and} \quad \lim_{x \rightarrow 0} (-x^2) = 0$$

Taking  $f(x) = -x^2$ ,  $g(x) = x^2 \sin(1/x)$ , and  $h(x) = x^2$  in the Squeeze Theorem, we obtain

$$\lim_{x \rightarrow 0} x^2 \sin \frac{1}{x} = 0$$




FIGURE 8  
 $y = x^2 \sin(1/x)$

## 1.6 Exercises



1. Given that

$$\lim_{x \rightarrow 2} f(x) = 4 \quad \lim_{x \rightarrow 2} g(x) = -2 \quad \lim_{x \rightarrow 2} h(x) = 0$$

find the limits that exist. If the limit does not exist, explain why.

- (a)  $\lim_{x \rightarrow 2} [f(x) + 5g(x)]$       (b)  $\lim_{x \rightarrow 2} [g(x)]^3$   
 (c)  $\lim_{x \rightarrow 2} \sqrt{f(x)}$       (d)  $\lim_{x \rightarrow 2} \frac{3f(x)}{g(x)}$   
 (e)  $\lim_{x \rightarrow 2} \frac{g(x)}{h(x)}$       (f)  $\lim_{x \rightarrow 2} \frac{g(x)h(x)}{f(x)}$

2. The graphs of  $f$  and  $g$  are given. Use them to evaluate each limit, if it exists. If the limit does not exist, explain why.



- (a)  $\lim_{x \rightarrow 2} [f(x) + g(x)]$       (b)  $\lim_{x \rightarrow 1} [f(x) + g(x)]$   
 (c)  $\lim_{x \rightarrow 0} [f(x)g(x)]$       (d)  $\lim_{x \rightarrow -1} \frac{f(x)}{g(x)}$   
 (e)  $\lim_{x \rightarrow 2} [x^3 f(x)]$       (f)  $\lim_{x \rightarrow -1} \sqrt{3 + f(x)}$

3–9 Evaluate the limit and justify each step by indicating the appropriate Limit Law(s).

3.  $\lim_{x \rightarrow -2} (3x^4 + 2x^2 - x + 1)$   
 4.  $\lim_{x \rightarrow -1} (x^4 - 3x)(x^2 + 5x + 3)$   
 5.  $\lim_{t \rightarrow -2} \frac{t^4 - 2}{2t^2 - 3t + 2}$       6.  $\lim_{u \rightarrow -2} \sqrt{u^4 + 3u + 6}$   
 7.  $\lim_{x \rightarrow 8} (1 + \sqrt[3]{x})(2 - 6x^2 + x^3)$       8.  $\lim_{t \rightarrow 2} \left( \frac{t^2 - 2}{t^3 - 3t + 5} \right)^2$   
 9.  $\lim_{x \rightarrow 2} \sqrt{\frac{2x^2 + 1}{3x - 2}}$

10. (a) What is wrong with the following equation?

$$\frac{x^2 + x - 6}{x - 2} = x + 3$$

(b) In view of part (a), explain why the equation

$$\lim_{x \rightarrow 2} \frac{x^2 + x - 6}{x - 2} = \lim_{x \rightarrow 2} (x + 3)$$

is correct.

11–32 Evaluate the limit, if it exists.

11.  $\lim_{x \rightarrow 2} \frac{x^2 + x - 6}{x - 2}$

12.  $\lim_{x \rightarrow -4} \frac{x^2 + 5x + 4}{x^2 + 3x - 4}$

13.  $\lim_{x \rightarrow 2} \frac{x^2 - x + 6}{x - 2}$

14.  $\lim_{x \rightarrow -1} \frac{x^2 - 4x}{x^2 - 3x - 4}$

15.  $\lim_{t \rightarrow -3} \frac{t^2 - 9}{2t^2 + 7t + 3}$

16.  $\lim_{x \rightarrow -1} \frac{2x^2 + 3x + 1}{x^2 - 2x - 3}$

17.  $\lim_{h \rightarrow 0} \frac{(-5 + h)^2 - 25}{h}$

18.  $\lim_{h \rightarrow 0} \frac{(2 + h)^3 - 8}{h}$

19.  $\lim_{x \rightarrow -2} \frac{x + 2}{x^3 + 8}$

20.  $\lim_{t \rightarrow 1} \frac{t^4 - 1}{t^3 - 1}$

21.  $\lim_{h \rightarrow 0} \frac{\sqrt{9 + h} - 3}{h}$

22.  $\lim_{u \rightarrow 2} \frac{\sqrt{4u + 1} - 3}{u - 2}$

23.  $\lim_{x \rightarrow -4} \frac{\frac{1}{4} + \frac{1}{x}}{4 + x}$

24.  $\lim_{x \rightarrow -1} \frac{x^2 + 2x + 1}{x^4 - 1}$

25.  $\lim_{t \rightarrow 0} \frac{\sqrt{1+t} - \sqrt{1-t}}{t}$

26.  $\lim_{t \rightarrow 0} \left( \frac{1}{t} - \frac{1}{t^2 + t} \right)$

27.  $\lim_{x \rightarrow 16} \frac{4 - \sqrt{x}}{16x - x^2}$

28.  $\lim_{h \rightarrow 0} \frac{(3 + h)^{-1} - 3^{-1}}{h}$

29.  $\lim_{t \rightarrow 0} \left( \frac{1}{t\sqrt{1+t}} - \frac{1}{t} \right)$

30.  $\lim_{x \rightarrow -4} \frac{\sqrt{x^2 + 9} - 5}{x + 4}$

31.  $\lim_{h \rightarrow 0} \frac{(x + h)^3 - x^3}{h}$

32.  $\lim_{h \rightarrow 0} \frac{\frac{1}{(x+h)^2} - \frac{1}{x^2}}{h}$

33. (a) Estimate the value of

$$\lim_{x \rightarrow 0} \frac{x}{\sqrt{1 + 3x} - 1}$$

by graphing the function  $f(x) = x/(\sqrt{1 + 3x} - 1)$ .(b) Make a table of values of  $f(x)$  for  $x$  close to 0 and guess the value of the limit.

(c) Use the Limit Laws to prove that your guess is correct.

34. (a) Use a graph of

$$f(x) = \frac{\sqrt{3+x} - \sqrt{3}}{x}$$

to estimate the value of  $\lim_{x \rightarrow 0} f(x)$  to two decimal places.(b) Use a table of values of  $f(x)$  to estimate the limit to four decimal places.

(c) Use the Limit Laws to find the exact value of the limit.

35. Use the Squeeze Theorem to show that  $\lim_{x \rightarrow 0} (x^2 \cos 20\pi x) = 0$ . Illustrate by graphing the functions  $f(x) = -x^2$ ,  $g(x) = x^2 \cos 20\pi x$ , and  $h(x) = x^2$  on the same screen.

36. Use the Squeeze Theorem to show that

$$\lim_{x \rightarrow 0} \sqrt{x^3 + x^2} \sin \frac{\pi}{x} = 0$$

Illustrate by graphing the functions  $f$ ,  $g$ , and  $h$  (in the notation of the Squeeze Theorem) on the same screen.37. If  $4x - 9 \leq f(x) \leq x^2 - 4x + 7$  for  $x \geq 0$ , find  $\lim_{x \rightarrow 4} f(x)$ .38. If  $2x \leq g(x) \leq x^4 - x^2 + 2$  for all  $x$ , evaluate  $\lim_{x \rightarrow 1} g(x)$ .39. Prove that  $\lim_{x \rightarrow 0} x^4 \cos \frac{2}{x} = 0$ .40. Prove that  $\lim_{x \rightarrow 0^+} \sqrt{x} [1 + \sin^2(2\pi/x)] = 0$ .

41–46 Find the limit, if it exists. If the limit does not exist, explain why.

41.  $\lim_{x \rightarrow 3} (2x + |x - 3|)$

42.  $\lim_{x \rightarrow -6} \frac{2x + 12}{|x + 6|}$

43.  $\lim_{x \rightarrow 0.5^-} \frac{2x - 1}{|2x^3 - x^2|}$

44.  $\lim_{x \rightarrow -2} \frac{2 - |x|}{2 + x}$

45.  $\lim_{x \rightarrow 0^-} \left( \frac{1}{x} - \frac{1}{|x|} \right)$

46.  $\lim_{x \rightarrow 0^+} \left( \frac{1}{x} - \frac{1}{|x|} \right)$

47. The *signum* (or *sign*) *function*, denoted by  $\text{sgn}$ , is defined by

$$\text{sgn } x = \begin{cases} -1 & \text{if } x < 0 \\ 0 & \text{if } x = 0 \\ 1 & \text{if } x > 0 \end{cases}$$

(a) Sketch the graph of this function.

(b) Find each of the following limits or explain why it does not exist.

(i)  $\lim_{x \rightarrow 0^+} \text{sgn } x$

(ii)  $\lim_{x \rightarrow 0^-} \text{sgn } x$

(iii)  $\lim_{x \rightarrow 0} \text{sgn } x$

(iv)  $\lim_{x \rightarrow 0} |\text{sgn } x|$

48. Let

$$f(x) = \begin{cases} x^2 + 1 & \text{if } x < 1 \\ (x - 2)^2 & \text{if } x \geq 1 \end{cases}$$

(a) Find  $\lim_{x \rightarrow 1^-} f(x)$  and  $\lim_{x \rightarrow 1^+} f(x)$ .(b) Does  $\lim_{x \rightarrow 1} f(x)$  exist?(c) Sketch the graph of  $f$ .

49. Let  $g(x) = \frac{x^2 + x - 6}{|x - 2|}$ .

(a) Find

(i)  $\lim_{x \rightarrow 2^+} g(x)$       (ii)  $\lim_{x \rightarrow 2^-} g(x)$

 (b) Does  $\lim_{x \rightarrow 2} g(x)$  exist?

 (c) Sketch the graph of  $g$ .

50. Let

$$g(x) = \begin{cases} x & \text{if } x < 1 \\ 3 & \text{if } x = 1 \\ 2 - x^2 & \text{if } 1 < x \leq 2 \\ x - 3 & \text{if } x > 2 \end{cases}$$

(a) Evaluate each of the following, if it exists.

(i)  $\lim_{x \rightarrow 1^-} g(x)$       (ii)  $\lim_{x \rightarrow 1} g(x)$       (iii)  $g(1)$

(iv)  $\lim_{x \rightarrow 2^-} g(x)$       (v)  $\lim_{x \rightarrow 2^+} g(x)$       (vi)  $\lim_{x \rightarrow 2} g(x)$

 (b) Sketch the graph of  $g$ .

 51. (a) If the symbol  $\llbracket x \rrbracket$  denotes the greatest integer function defined in Example 10, evaluate

(i)  $\lim_{x \rightarrow -2^+} \llbracket x \rrbracket$       (ii)  $\lim_{x \rightarrow -2} \llbracket x \rrbracket$       (iii)  $\lim_{x \rightarrow -2.4} \llbracket x \rrbracket$

 (b) If  $n$  is an integer, evaluate

(i)  $\lim_{x \rightarrow n^-} \llbracket x \rrbracket$       (ii)  $\lim_{x \rightarrow n^+} \llbracket x \rrbracket$

 (c) For what values of  $a$  does  $\lim_{x \rightarrow a} \llbracket x \rrbracket$  exist?

 52. Let  $f(x) = \llbracket \cos x \rrbracket$ ,  $-\pi \leq x \leq \pi$ .

 (a) Sketch the graph of  $f$ .

(b) Evaluate each limit, if it exists.

(i)  $\lim_{x \rightarrow 0} f(x)$       (ii)  $\lim_{x \rightarrow (\pi/2)^-} f(x)$

(iii)  $\lim_{x \rightarrow (\pi/2)^+} f(x)$       (iv)  $\lim_{x \rightarrow \pi/2} f(x)$

 (c) For what values of  $a$  does  $\lim_{x \rightarrow a} f(x)$  exist?

 53. If  $f(x) = \llbracket x \rrbracket + \llbracket -x \rrbracket$ , show that  $\lim_{x \rightarrow 2} f(x)$  exists but is not equal to  $f(2)$ .

54. In the theory of relativity, the Lorentz contraction formula

$$L = L_0 \sqrt{1 - v^2/c^2}$$

expresses the length  $L$  of an object as a function of its velocity  $v$  with respect to an observer, where  $L_0$  is the length of the object at rest and  $c$  is the speed of light. Find  $\lim_{v \rightarrow c^-} L$  and interpret the result. Why is a left-hand limit necessary?

 55. If  $p$  is a polynomial, show that  $\lim_{x \rightarrow a} p(x) = p(a)$ .

 56. If  $r$  is a rational function, use Exercise 55 to show that  $\lim_{x \rightarrow a} r(x) = r(a)$  for every number  $a$  in the domain of  $r$ .

 57. If  $\lim_{x \rightarrow 1} \frac{f(x) - 8}{x - 1} = 10$ , find  $\lim_{x \rightarrow 1} f(x)$ .

 58. If  $\lim_{x \rightarrow 0} \frac{f(x)}{x^2} = 5$ , find the following limits.

(a)  $\lim_{x \rightarrow 0} f(x)$       (b)  $\lim_{x \rightarrow 0} \frac{f(x)}{x}$

59. If

$$f(x) = \begin{cases} x^2 & \text{if } x \text{ is rational} \\ 0 & \text{if } x \text{ is irrational} \end{cases}$$

 prove that  $\lim_{x \rightarrow 0} f(x) = 0$ .

 60. Show by means of an example that  $\lim_{x \rightarrow a} [f(x) + g(x)]$  may exist even though neither  $\lim_{x \rightarrow a} f(x)$  nor  $\lim_{x \rightarrow a} g(x)$  exists.

 61. Show by means of an example that  $\lim_{x \rightarrow a} [f(x)g(x)]$  may exist even though neither  $\lim_{x \rightarrow a} f(x)$  nor  $\lim_{x \rightarrow a} g(x)$  exists.

 62. Evaluate  $\lim_{x \rightarrow 2} \frac{\sqrt{6-x} - 2}{\sqrt{3-x} - 1}$ .

 63. Is there a number  $a$  such that

$$\lim_{x \rightarrow -2} \frac{3x^2 + ax + a + 3}{x^2 + x - 2}$$

 exists? If so, find the value of  $a$  and the value of the limit.

 64. The figure shows a fixed circle  $C_1$  with equation  $(x - 1)^2 + y^2 = 1$  and a shrinking circle  $C_2$  with radius  $r$  and center the origin.  $P$  is the point  $(0, r)$ ,  $Q$  is the upper point of intersection of the two circles, and  $R$  is the point of intersection of the line  $PQ$  and the  $x$ -axis. What happens to  $R$  as  $C_2$  shrinks, that is, as  $r \rightarrow 0^+$ ?
